

⑨



Europäisches Patentamt
European Patent Office
Office européen des brevets

⑩ Publication number:

0 369 583
A1

⑫

EUROPEAN PATENT APPLICATION

⑪ Application number: 89309463.1

⑤ Int. Cl.⁵ C07D 473/32, C07D 473/40,
A61K 31/52

⑬ Date of filing: 18.09.89

Claims for the following Contracting State: ES.

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

③① Priority: 21.09.88 GB 8822236

④③ Date of publication of application:
23.05.90 Bulletin 90/21

③④ Designated Contracting States:
BE CH DE ES FR GB GR IT LI NL

⑦① Applicant: BEECHAM GROUP PLC
Beecham House Great West Road
Brentford Middlesex TW8 9BD(GB)

⑦② Inventor: Geen, Graham Richard Beecham
Pharmaceuticals


Great Burgh Yew Tree Bottom Road
Epsom Surrey KT18 5XQ(GB)

Inventor: Hanson, John Christopher Beecham
Pharmaceuticals
Brockham Park
Betchworth Surrey RH3 7AJ(GB)

⑦④ Representative: Jones, Pauline et al
Beecham Pharmaceuticals Patents & Trade
Marks Dept. Great Burgh Yew Tree Bottom
Road
Epsom Surrey KT18 5XQ(GB)

⑤④ Chemical process for the preparation of purine derivatives.

⑤⑦ A process for the preparation of a compound of formula (I)




(I)

which process comprises reacting a compound of formula (II):


EP 0 369 583 A1

REFERENCE BA



(II)

wherein the amino group is optionally protected, with a side chain intermediate of formula (III):




(III)

wherein Q is a leaving group, R_x and R_y are protected hydroxymethyl or acyloxymethyl, or group(s) convertible to hydroxymethyl or acyloxymethyl; and R_z is hydrogen or a group convertible thereto; and thereafter converting the 6- and 8- chloro substituents to hydrogen by means of reduction; converting R_x and R_y when other than hydroxymethyl or acyloxymethyl, to hydroxymethyl or acyloxymethyl, optionally converting R_x, R_y hydroxymethyl to acyloxymethyl or vice versa, deprotecting the 2-amino group where necessary and converting R_z , (when other than hydrogen) to hydrogen; and optionally forming a pharmaceutically acceptable salt thereof.

CHEMICAL PROCESS

The present invention relates to a novel chemical process for the preparation of purine derivatives which have antiviral activity.


EP-A-182024 (Beecham Group p.l.c.) describes compounds of formula (I):



(I)

wherein R_a and R_b are independently hydrogen or a group $\text{RCO}-$ wherein R is phenyl or C_1-18 alkyl.


The compound of formula (B) within formula (I):



(B)


is of particular interest as an antiviral agent (BRL 42810 or famciclovir).

The process already described for the preparation of the compounds of formula (I) involves the reaction of a purine intermediate, 2-amino-6-chloropurine of formula (C):



(C)

with a side chain intermediate of formula (D):




(D)

wherein R_2 and R_3 are independently acyl groups or hydroxy protecting groups and Z is a leaving group, such as halo, for example chloro, bromo, iodo; and thereafter converting the 6-chloro group to hydrogen by reduction methods.

The intermediate of formula (C) is prepared by chlorination of guanine, as described in EP-A-203685 (Beecham Group p.l.c.). Despite this improvement in the method of production of 2-amino-6-chloropurine, the isolable yield at this chlorination step is sufficiently low that there is a need for alternative routes to compounds of formula (I).


It has surprisingly been discovered that, by introducing an 8-chloro substituent into the compound of formula (C), the yield of corresponding purine intermediate is increased, and therefore increasing the overall yield of the resulting compound of formula (I).

Accordingly, the present invention provides a process for the preparation of a compound of formula (I), or a pharmaceutically acceptable salt thereof, which process comprises reacting a purine intermediate of formula (II):



(II)


wherein the amino group is optionally protected, with a side chain intermediate of formula (III):



(III)

wherein Q is a leaving group, R_x and R_y are protected hydroxymethyl or acyloxymethyl, or group(s) convertible to hydroxymethyl or acyloxymethyl; and R_z is hydrogen or a group convertible thereto; and thereafter converting the 6- and 8- chloro substituents to hydrogen by means of reduction; converting R_x and R_y when other than hydroxymethyl or acyloxymethyl, to hydroxymethyl or acyloxymethyl, optionally converting R_x/R_y hydroxymethyl to acyloxymethyl or *vice versa*, deprotecting the 2-amino group where necessary and converting R_z , (when other than hydrogen) to hydrogen; and optionally forming a pharmaceutically acceptable salt thereof.

The intermediates formed in this reaction are of formula (IV):



(IV)

which are novel and form an aspect of the invention.

The reaction may be carried out in an inert solvent, for example dimethylformamide, dimethylsulphoxide or acetonitrile, preferably dimethylformamide, in the presence of an inorganic or organic base, over a temperature range from 0°C to the boiling point of the solvent, usually 30-40°C. Examples of inorganic bases include alkali metal hydrides, alkali metal carbonates such as sodium or potassium carbonate and preferably potassium carbonate. Suitable organic bases are 1,8-diazabicyclo[5.4.0]undec-7-ene and tetramethyl guanidine.

Suitable examples of the leaving group Q, include halo, such as chloro, bromo or iodo, and tosyloxy and mesyloxy.

Suitable examples of hydroxy protecting groups (other than acyl groups) include the *t*-Butyl dimethylsilyl group removable by 80% acetic acid at elevated temperatures, around 90°C, or by treatment with tetrabutyl ammonium fluoride in a solvent, such as tetrahydrofuran, at ambient temperature.

Another suitable protecting group is wherein the two hydroxy groups in formula (III) (when R_x is hydroxymethyl) are reacted with 2,2-dimethoxypropane, forming a 1,3-dioxan ring. This group may be removed by acidic hydrolysis.


Other suitable protecting groups include substituted benzyl groups such as *p*-methoxybenzyl, removable by treatment with 2,3-dichloro-5,6-dicyanobenzoquinone.

Other suitable protecting groups are apparent to those skilled in the art.


R_x and/or R_y may be acyloxymethyl, such as a group RCO₂CH₂ wherein R is as defined in formula (I). Examples of R include methyl, ethyl, *n*- and *iso*-propyl, *n*- and *iso*-, *sec*- and *tert*-butyl, preferably methyl.

Interconversion of R_x/R_y acyloxymethyl and hydroxymethyl may be carried out conventionally as described in EP-A-141927.

Other suitable values of R_x, R_y, R_z include wherein the compounds of formula (III) is of formula (IIIA) or (IIIB)




(IIIA)




(IIIB)

wherein R_p and R_q are independently hydrogen, C₁₋₅ alkyl or phenyl, or R_p and R_q together are C₄₋₆ polymethylene; and R_r is C₁₋₆ alkyl or phenyl C₁₋₆ alkyl, in which any phenyl moieties are optionally substituted. Optional substituents include one or two groups selected from C₁₋₄ alkyl, halo and C₁₋₄ alkoxy. Halo includes iodo, bromo, chloro and fluoro; alkoxy/alkyl includes those containing methyl, ethyl, *n*- and *iso*-propyl.

When the compound of formula (IIIA) is used, the resulting intermediate is of formula (IVA):




When the compound of formula (IIIB) is used, the resulting intermediate is of formula (IVB):




Values for alkyl groups R_p and R_q and R_r include these values listed as suitable for R in formula (I), preferably methyl for R_p and R_q and ethyl for R_r . In addition R_p and R_q may together be C_4 or C_5 polymethylene.

The intermediates of formulae (IVA) and (IVB) are subsequently converted to an intermediate of formula (V):




by transesterification and hydrolysis/decarboxylation respectively by methods analogous to those disclosed in EP-A-0302644.

An intermediate of formula (V) is convertible to a compound of formula (VI):



by reduction, under conventional conditions using, for example, sodium borohydride.

It is preferred, however, that the intermediate of formula (III) is of formula (III)':



for the preparation of compounds of formula (A) and (B) as defined, because:

i) Compounds of formula (III)' give a particularly good N9:N7 ratio (regioselectivity).

ii) Ease of separation of N9:N7 isomers.

The 2-amino group may be protected, for example, using a benzyl protecting group, removable by hydrogenolysis. It may also be protected by an acyl group, for example acetyl, removable by hydrolysis.


The conversion of the chloro substituents to hydrogen by means of reduction takes place conventionally, according to the methods described for reduction of the 6-chloro substituent in EP-A-182024.

Pharmaceutically acceptable salts are formed conventionally.

The purine intermediate of formula (II) is novel and forms an aspect of the present invention.

It may be prepared by chlorination of guanine by:

i) reacting guanine with a sulphur oxychloride in the presence of chloride ions, to give 8-chloroguanine of formula (VII):



Suitable sulphur oxychlorides include SOCl_2 , SO_2Cl_2 , MeSO_3Cl and TsCl . Suitable sources of chloride ions include quaternary ammonium chlorides such as tetraalkyl substituted ammonium chlorides, for example Et_4NCl , Et_3MeNCl , $\text{Et}_2\text{Me}_2\text{NCl}$, EtMe piperidinium chloride and Et_2NPh . HCl . The reaction takes place at elevated temperatures $30-100^\circ\text{C}$ preferably $50-80^\circ\text{C}$ in a solvent such as acetonitrile, dichloromethane, chloroform or methanesulphonic acid. Alternatively, the chlorinating agent may also function as the solvent.

ii) reacting 8-chloroguanine with phosphorus oxychloride in the presence of chloride ions, in analogous manner to the chlorination of guanine itself, described in EP-A-203685.

It will be appreciated that steps i) and ii) may be carried out in reverse order, via 2-amino-6-chloropurine, although this route is less preferred.

Intermediates of formula (III) wherein R_x/R_y are protected hydroxymethyl or acyloxymethyl may be prepared as described in EP-A-141927 and EP-A-182024 (both Beecham Group p.l.c), or by analogous methods thereto.

Intermediates of the formula (IIIA) are known or are prepared by analogous methods, such as that

described in Organic Syntheses Vol 60, page 66.


Intermediates of formula (IIIB) are known or prepared by analogous methods. The compound of formula (IIIB) wherein Q is bromo and R, is ethyl may be prepared from triethyl methanetricarboxylate according to the procedure described by H. Rapoport et al., J. Org. Chem., 44, 3492(1979).

It will also be appreciated that the side chain intermediate of formula (III) may be replaced by an appropriate side chain for the preparation of other 6-deoxy guanine antiviral agents, such as those described in EP-A-108285 (The Wellcome Foundation Limited) and EP-A-186640 (Astra Läkemedel Aktiebolag).

The following Examples illustrate the invention.

Description

Preparation of 2-amino-8-chloro-6-hydroxypurine (8-chloroguanine) and its hydrochloride salt.



Method 1

A mixture of guanine (22.5g), methyltriethylammonium chloride (88g) and thionyl chloride (50ml) was slowly heated and stirred at 50-70°C for ½h, then kept at 70°C for a further ½h. The reaction mixture was then cooled and the excess thionyl chloride removed under vacuum. The residue was poured into water and the pH adjusted to 7 to give a tan-coloured precipitate. In order to expedite filtration and to achieve further purification, the pH was further adjusted to 13 with 40% sodium hydroxide solution, any insoluble material filtered off, and the resulting solution acidified slowly to pH 8 allowing the product to crystallize, particularly at pH 11.5. The solid was filtered washed with acetone and air dried at 40°C to give the title compound as a buff solid (28.5g) (water assay 14.1%). Vacuum drying at 40°C afforded a solid analysing as a monohydrate.

I.R. 3380, 3200, 3100, 1670 cm⁻¹.

M.S. (negative ion F.A.B.) 184, 186 (M-H)

Found: C, 29.33; H, 2.62; N, 34.59%. C₅H₄N₅OCl.H₂O requires: C, 29.50; H, 2.97; N, 34.40%.

Method 2

Sulphuryl chloride (3.3ml) was added to a stirred mixture of guanine (3.02g) and methyltriethylammonium chloride (13g) in acetonitrile (22ml). The reaction temperature increased slowly to 50°C, and was maintained at 50°C for a further 45 minutes by heating. The mixture was then cooled, poured into water and the pH adjusted to 7. The crude product was isolated by filtration or purified as in Method 1.

Method 3

Guanine (1.52g) was dissolved with warming in a mixture of methane sulphonic acid (10ml), thionyl chloride (10ml), and tetraethylammonium chloride (6g). The mixture was heated to 80°C for 1h, cooled and poured into water. The product was isolated as in Method 1. (Thionyl chloride may be replaced by phosphorus oxychloride or methanesulphonyl chloride).


HCl salt

8-chloroguanine (2g), dissolved in hot water (40ml) and concentrated hydrochloric acid (25ml), was treated with charcoal, filtered and the filtrate cooled to 4 °C overnight to give the title compound as an off-white solid (1.2g) analysing as a monohydrate.

Found: C, 24.73; H, 3.25; N, 29.36; Cl, 28.95%. $C_5H_4N_5OCl \cdot HCl \cdot H_2O$

requires: C, 25.02; H, 2.94; N, 29.18; Cl 29.54%.

^{13}C n.m.r. (D_6 -DMSO) 112.0, 135.6, 149.4 152.7, 152.9 p.p.m.

Description 2Preparation of 2-Amino-6,8-dichloropurineMethod 1

8-chloroguanine (vacuum dried at 115 °C over phosphorus pentoxide) (2.04g) was added to a solution of methyl triethylammonium chloride (6.6g) in acetonitrile (11ml). Phosphorus oxychloride (5.6ml) was then added and the mixture heated to 60 °C for 1h. The mixture was then cooled, and the solvent and excess phosphorus oxychloride removed under vacuum. Water was added, the pH adjusted to 3 and the mixture stirred overnight at room temperature to complete hydrolysis of the phosphorus containing intermediates. The pH was then adjusted to 10 the solution treated with charcoal, filtered and acidified slowly to pH 5. The fine crystals were filtered off, washed with acetone and air dried at 40 °C, giving the title compound as an off-white solid (1.78g), analysing as a hemihydrate (water assay 4.1%).

I.R. 3330, 3190, 1680, 1630, 1570 cm^{-1} .

M.S. (electron impact) 203, 205, 207 M^+ (Cl_2 isotope pattern); 168, 170 ($M^+ - Cl$)

^{13}C n.m.r. (D_6 -DMSO) 122.9, 138.6, 147.4, 155.9, 159.8 p.p.m.

Found: C, 28.29; H, 1.92; N, 33.27%. $C_5H_3N_5Cl_2 \cdot \frac{1}{2}H_2O$


requires: C, 28.19; H, 1.89; N, 32.88%.

U.V. (EtOH) λ_{max} 219, 246, 311 nm.

Method 2

2-Amino-6-chloropurine (3.4g) was added to a solution of methyl triethyl ammonium chloride (13g) in acetonitrile (22ml). Sulphuryl chloride (5ml) was added and the reaction mixture heated to 60 °C for 24h, cooled, poured into water (150ml) and the pH adjusted to 9 with 40% sodium hydroxide solution. Charcoal was then added, the mixture filtered and the filtrate acidified slowly to pH 4.5 to give the title compound as a solid which was filtered off (3.9g).

Example9-(4-Acetoxy-3-acetoxymethylbut-1-yl)-2-aminopurine (BRL 42810)



Method 1

(a) 2-Acetoxymethyl-4-iodobutyl-1-acetate (9.4g) was added to a stirred mixture of 2-amino-6,8-dichloropurine (5.8g) and anhydrous potassium carbonate (5.9g) in N,N-dimethylformamide (100ml), and the resulting mixture stirred at ambient temperature overnight. The reaction mixture was filtered and the filtrate evaporated to give a thick gum which was shown by ^1H n.m.r. analysis to be a mixture of the N-9 and N-7 alkylated purines in the ratio 17:1 respectively. T.l.c. (5% methanol-dichloromethane) showed two products of $r_f = 0.24, 0.32$; corresponding to the N-7 and N-9 alkylated purines. Column chromatography on silica (eluant 2.5% methanol-dichloromethane) of the gum afforded 9-(4-acetoxy-3-acetoxymethylbut-1-yl)-2-amino-6,8-dichloropurine, which was crystallised from ethyl acetate-diethyl ether to give a colourless solid (4.7g) m.p. $118-119^\circ\text{C}$.

^1H n.m.r. (D^6 -DMSO) δ 1.83 (q, 2H, $>\text{CHCH}_2\text{CH}_2-$), 2.03 (s, 6H, $-\text{CH}_3$), 2.05 (m, 1H, $>\text{CH}-$), 4.07 (m, 4H, $-\text{CH}_2\text{O}-$), 4.15 (t, 2H, $>\text{NCH}_2-$), 7.05 (brs, 2H, $-\text{NH}_2$).

Found: C, 43.13; H, 4.48; N, 18.07%. $\text{C}_{14}\text{H}_{17}\text{N}_5\text{O}_4\text{Cl}_2$ requires: C, 43.09; H, 4.39; N, 17.95%.

(b) A mixture of 9-(4-acetoxy-3-acetoxymethylbut-1-yl)-2-amino-6,8-dichloropurine (3.9g), 5% palladium on charcoal (0.5g) and triethylamine (3.1ml) was suspended in ethanol (250ml) and hydrogenated at 50 p.s.i. overnight. The reaction was initially heated to 60°C , and then allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated. The residue was taken up in water (100ml) and extracted with dichloromethane (2x200ml). The organic extracts were combined and dried over magnesium sulphate. Filtration and evaporation gave the title compound as a gum which crystallized on standing. Recrystallization from butan-1-ol gave a colourless solid (2.4g) m.p. $102-103^\circ\text{C}$.

^1H n.m.r. (D^6 -DMSO) δ 1.88 (m, 3H, $>\text{CHCH}_2\text{CH}_2-$), 2.00 (s, 6H, $-\text{CH}_3$), 4.03 (d, 4H, $-\text{CH}_2\text{O}-$), 4.14 (t, 2H, $>\text{NCH}_2-$), 6.45 (brs, 2H, $-\text{NH}_2$), 8.09 (s, 1H, H-8), 8.57 (s, 1H, H-6).

Method 2

A mixture of 2-amino-6,8-dichloropurine (2.04g), 6,6-dimethyl-5,7-dioxaspiro[2.5]octane-4,8-dione (1.73g) and anhydrous potassium carbonate (2.12g) in N,N-dimethylformamide (40ml) was stirred at room temperature for 18 hours under dry nitrogen. The solvent was evaporated and the residue dissolved in water (60ml) and acidified from pH 10 to pH 4 with dilute hydrochloric acid. The yellow precipitate was filtered off, washed with water and dried to give 2-amino-6,8-dichloro-9-[1-(2,2-dimethyl-1,3-dioxane-4,6-dione-5-yl)ethyl]-purine (2.2g, 50%). T.l.c. r_f 0.65 (25% methanol-dichloromethane).

^1H n.m.r. (D^6 -DMSO) δ 1.69 (s, 3H, CH_3), 1.79 (s, 3H, CH_3), 2.42 (m, 2H, CHCH_2), 4.31 (t, 2H, NCH_2), 4.4 (vbr s, 2H, $\text{NH}_2 + \text{H}_2\text{O}$), 4.56 (t, 1H, CHCH_2).

BRL 42810 is prepared from the above compound by methods analogous to those disclosed in EP-A-302644:-

- i) Reduction (see Description 12)
- ii) Trans-esterification (see Example 2)
- iii) Reduction (see Method d))
- iv) Acetylation (see Method c))

Method 3

A mixture of 2-amino-6,8-dichloropurine (0.82g), 5-bromoethyl-2,2-dimethyl-1,3-dioxan (0.9g) and anhy-

drous potassium carbonate (0.83g) in N,N-dimethylformamide (20ml) was stirred at room temperature for 18 hours under dry nitrogen. The mixture was then filtered and the filtrate evaporated. The residue was purified by column chromatography on silica (100g) [eluent 2% methanol-dichloromethane] to give 2-amino-6,8-dichloro-9-(2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl)purine (t.l.c. r.f. 0.50, 5% methanol-dichloromethane) as a pale solid (0.62g, 45%). Recrystallisation from diethyl ether gave an off-white solid m.p. 149-150 °C.

¹H n.m.r. (D⁶-DMSO): δ 1.28(s,3H,CH₃), 1.33(s,3H,CH₃), 1.62(m,1H,CHCH₂), 1.70(m,2H,CH₂CH₂CH), 3.56-(dd,2H,2XH_{ax}), 3.84(dd,2H,2XH_{eq}), 4.06(t,2H, NCH₂), 7.06(br s,2H,NH₂).

Elemental analysis: Found: C: 45.01, H: 4.89, N:20.25; C₁₃H₁₇N₅O₂Cl₂ requires: C: 45.10, H: 4.95, N:20.23.

BRL 42810 is prepared from the above compound by methods analogous to those disclosed in EP-A-

302644:-

- i) Reduction (see Description 12)
- ii) Hydrolysis (see Method e))
- iii) Acetylation (see Method c))

Method 4

Ethyl 4-bromo-2,2-dicarboethoxybutanoate (8.2g) was added to a stirred mixture of 2-amino-6,8-dichloropurine (4.08g) and anhydrous potassium carbonate (4.14g) in N,N-dimethylformamide (80 cm³) and the resulting mixture stirred at 70-80 °C for 72 hours under dry nitrogen. The reaction mixture was cooled to room temperature, filtered and the filtrate evaporated. T.l.c. [3% methanol-dichloromethane] showed one major spot r.f. = 0.44. The residue was purified by column chromatography on silica (200g) [eluent = 2% methanol-dichloromethane] to give 2-amino-6,8-dichloro-9-(ethyl-2,2-dicarboethoxybutanoate-4-yl) purine (2.6g 28%) as a yellow solid. Recrystallization from ethyl acetate/diethyl ether gave an off-white solid m.p. 153-155 °C.

¹H n.m.r. (D⁶-DMSO) δ 1.22(t,9H,-CH₂CH₃), 2.50(t,2H,-CH₂C), 4.23(q,6H,-CH₂CH₃), 4.31(t,2H, N-CH₂), 7.02 (br s,2H,NH₂).


Elemental analysis: Found: C: 44.35, H: 4.57, N: 15.09; C₁₇H₂₁N₅O₆Cl₂ requires: C: 44.17, H: 4.58, N: 15.15.

BRL 42810 is prepared from the above compound by methods analogous to those disclosed in EP-A-302644:-


- i) Reduction (see Description 12)
- ii) Decarbethoxylation (see Example 3)
- iii) Reduction (see Method b))
- iv) Acylation (see Method c))

Claims

1. A process for the preparation of a compound of formula (I)




which process comprises reacting a compound of formula (II):



(II)


wherein the amino group is optionally protected, with a side chain intermediate of formula (III):




(III)

wherein Q is a leaving group, R_x and R_y are protected hydroxymethyl or acyloxymethyl, or group(s) convertible to hydroxymethyl or acyloxymethyl; and R_z is hydrogen or a group convertible thereto; and thereafter converting the 6- and 8- chloro substituents to hydrogen by means of reduction; converting R_x and R_y when other than hydroxymethyl or acyloxymethyl, to hydroxymethyl or acyloxymethyl, optionally converting R_x , R_y hydroxymethyl to acyloxymethyl or vice versa, deprotecting the 2-amino group where necessary and converting R_z , (when other than hydrogen) to hydrogen; and optionally forming a pharmaceutically acceptable salt thereof.

2. A process according to claim 1 wherein the compound of formula (III) is of formula (IIIA) or (IIIB):




(IIIA)



(IIIB)

wherein R_p and R_q are independently hydrogen, C_{1-6} alkyl or phenyl, or R_p and R_q together are C_{4-6} polymethylene; and R_r is C_{1-6} alkyl or phenyl C_{1-6} alkyl, in which any phenyl moieties are optionally substituted.

3. A process according to claim 1 wherein the compound of formula (III) is of formula (III)':




(III)'


wherein Q is as defined in claim 1.

4. A process as defined in any one of the claims 1 to 3 wherein Q is fluoro, chloro, bromo, iodo, tosyloxy or mesyloxy.

5. A process according to any one of the claims 1 to 4 for the preparation of a compound of formula (IB):



6. A compound of formula (IV) as defined in claim 1:




7. 9-(4-Acetoxy-3-acetoxymethylbutyl-1-yl)-2-amino-6,8-dichloropurine,
2-Amino-6,8-dichloro-9-[1-(2,2-dimethyl-1,3-dioxane-4,6-dione-5-yl)eth-2-yl]purine,
2-Amino-6,8-dichloro-9(2-(2,2-dimethyl-1,3-dioxan-5-yl)ethyl)purine, or
2-Amino-6,8-dichloro-9-(ethyl-2,2-dicarboethoxybutanoate-4-yl)purine.

8. 2-Amino-6,8-dichloropurine.


9. A process for the preparation of a 9-N-substituted 6-deoxyguanine antiviral agent, which process comprises reacting a compound of formula (II) as defined in claim 1, with an appropriate side chain; and thereafter converting the 6- and 8- chloro substituents to hydrogen by means of reduction, deprotecting the 2-amino group where necessary.

Claims for the following Contracting State : ES

1. A process for the preparation of a compound of formula (I)




which process comprises reacting a compound of formula (II):



(II)


wherein the amino group is optionally protected, with a side chain intermediate of formula (III):




(III)

wherein Q is a leaving group, R_x and R_y are protected hydroxymethyl or acyloxymethyl, or group(s) convertible to hydroxymethyl or acyloxymethyl; and R_z is hydrogen or a group convertible thereto; and thereafter converting the 6- and 8- chloro substituents to hydrogen by means of reduction; converting R_x and R_y when other than hydroxymethyl or acyloxymethyl, to hydroxymethyl or acyloxymethyl, optionally converting $R_x R_y$ hydroxymethyl to acyloxymethyl or vice versa, deprotecting the 2-amino group where necessary and converting R_z , (when other than hydrogen) to hydrogen; and optionally forming a pharmaceutically acceptable salt thereof.

2. A process according to claim 1 wherein the compound of formula (III) is of formula (IIIA) or (IIIB):




(IIIA)



(IIIB)

wherein R_p and R_q are independently hydrogen, C_{1-5} alkyl or phenyl, or R_p and R_q together are C_{4-5} polymethylene; and R_r is C_{1-5} alkyl or phenyl C_{1-5} alkyl, in which any phenyl moieties are optionally substituted.

3. A process according to claim 1 wherein the compound of formula (III) is of formula (III)':




(III)'


wherein Q is as defined in claim 1.

4. A process as defined in any one of the claims 1 to 3 wherein Q is fluoro, chloro, bromo, iodo, tosyloxy or mesyloxy.

5. A process according to any one of the claims 1 to 4 for the preparation of a compound of formula (IB):



6. A process according to any one of claims 1, 3, 4 or 5 for the preparation of a compound of formula (IV) as defined in claim 1:



7. A process for the preparation of an intermediate compound produced by reacting a compound of formula (II) with a compound of formula (III) as defined in claim 1, wherein the intermediate compound is selected from:

- 30
- 9-(4-Acetoxy-3-acetoxymethylbutyl-1-yl)-2-amino-6,8-dichloropurine,
 - 2-Amino-6,8-dichloro-9-[1-(2,2-dimethyl-1,3-dioxane-4,6-dione-5-yl)eth-2-yl]purine,
 - 2-Amino-6,8-dichloro-9-(2,2-dimethyl-1,3-dioxan-5-yl)ethylpurine, and
 - 35 2-Amino-6,8-dichloro-9-(ethyl-2,2-dicarboethoxybutanoate-4-yl)purine.

8. A process for the preparation of a 9-N-substituted 6-deoxyguanine antiviral agent, which process comprises reacting a compound of formula (II) as defined in claim 1, with an appropriate side chain; and thereafter converting the 6- and 8- chloro substituents to hydrogen by means of reduction, deprotecting the 2-amino group where necessary.

40

This Page Blank (uspto)



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 89309463 1
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D, A	<u>EP - A2 - 0 182 024</u> (BEECHAM) * Claims; pages 5-10 * --	1, 3-6, 9	C 07 D 473/32 C 07 D 473/40 A 61 K 31/52
A	<u>EP - A2 - 0 203 685</u> (BEECHAM) * Formula 1 * -----	8	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 D 473/00
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
VIENNA		07-12-1989	HOFBAUER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

This Page Blank (uspto)